

MYCOTAXON

<http://dx.doi.org/10.5248/127.25>

Volume 127, pp. 25–31

January–March 2014

Three new species of *Septobasidium* from Yunnan and Guangxi in ChinaWEI LI^{1, 2} & LIN GUO^{1*}¹State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China²University of Chinese Academy of Sciences, Beijing 100049, ChinaCORRESPONDENCE TO *: guol@im.ac.cn

ABSTRACT — Three new species are described: *Septobasidium brunneum* on *Eurya* sp. associated with *Chionaspis* sp., *Septobasidium guangxiense* associated with *Pseudaulacaspis* sp., and *Septobasidium transversum* associated with tribe *Lepidosaphidini*. They were collected from Yunnan Province and Guangxi Zhuang Autonomous Region in China.

KEY WORDS — *Pucciniomycetes*, *Septobasidiales*, *Septobasidiaceae*, taxonomy

Previously, 26 species of *Septobasidium* have been recorded in Yunnan and Guangxi of China. An additional three new species of *Septobasidium*, collected from these regions, are described as follows:

***Septobasidium brunneum* Wei Li bis & L. Guo, sp. nov.**

FIGS 1–7





FUNGAL NAME FN 570073

Differs from *Septobasidium thwaitesii* by its thinner section, smaller basidia, and absence of a horizontal section layer.

TYPE: China, Yunnan, Dali, Cangshan, on *Eurya* sp. (*Pentaphylacaceae*), associated with *Chionaspis* sp. (*Diaspididae*), 11.VIII.2011, F. He YN45 (HMAS 243152, **holotype**).

ETYMOLOGY: The epithet refers to the surface color of basidiomata.

Basidiomata on branches, resupinate, 7–9 cm long, 2–3 cm wide, dark brown or purple-brown; surface with many cracks, frequently extending out to form lateral margins in old stage, margin indeterminate. In section 350–700 µm thick. Subiculum brown, 30–50 µm thick. Pillars brown, 50–80 µm high, 80–150 µm wide. Hyphal layer brown, 150–220 µm high. Hymenium brown, 70–100 µm thick. The hyphae renew grown to form successive hyphal layers and hymenium from hymenial layer in old stage. Probasidia pyriform or






FIG 1. *Septobasidium brunneum* (HMAS 243152, holotype). Basidia.

subglobose, $10\text{--}15 \times 7\text{--}9\ \mu\text{m}$, persistent. Basidia cylindrical, curved or straight, 4-celled, $20\text{--}30 \times 4\text{--}6\ \mu\text{m}$, brown. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

COMMENTS: *Septobasidium brunneum* is similar to *S. thwaitesii* (Berk. & Broome) Pat., which differs by having a thicker section ($270\text{--}1500\ \mu\text{m}$), two distinct horizontal layers, and larger basidia ($37\text{--}46 \times 12\text{--}13.5\ \mu\text{m}$; Couch 1938).

***Septobasidium guangxiense* Wei Li bis & L. Guo, sp. nov.**

Figs 8–11

FUNGAL NAME FN 570074

Differs from *Septobasidium ligustri* by its numerous fissures on the basidiome surface, its larger basidia, and its horizontal section layer.

TYPE: China, Guangxi, Wuming, Damingshan Scenic Area, alt. 700 m, on unidentified plant, associated with *Pseudaulacaspis* sp. (*Diaspididae*), 29.VIII.2011, W. Li & S.H. He 1519 (HMAS 244542, holotype).

ETYMOLOGY: The epithet refers to the type locality, Guangxi.

Basidiomata on branches, resupinate, 40–50 cm long, 6–7 cm wide, yellowish brown or brown; surface with numerous fissures at maturity; margin determinate. In section 200–500 μm thick. Subiculum brown, 30–70 μm thick. Pillars brown, 70–100 μm high, 50–80 μm wide, forming a horizontal layer 20–40 μm thick on the top of pillars. Hyphal layer brown, 150–250 μm thick.

FIGS 2–7. *Septobasidium brunneum* (HMAS 243152, holotype). 2. Basidiomata on branches; 3–4. Sections of basidiomata; 5–6. Basidia (arrows); 7. Haustoria.




FIG 8. *Septobasidium guangxiense* (HMAS 244542, holotype). Basidia.

Hymenium brown, 50–100 μm thick, with irregularly arranged hyphae. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or curved, 4-celled, 27–38 $\times$ 5–10 μm , hyaline or brownish. Basidiospores not seen. Haustoria not seen.

COMMENTS: *Septobasidium guangxiense* is similar to *S. ligustri* C.X. Lu & L. Guo, which differs by having fewer fissures on the basidiome surface, lacking a horizontal section layer, and having smaller basidia (15–29 $\times$ 5–7.5 μm ; Lu & Guo 2010c).

***Septobasidium transversum* Wei Li bis & L. Guo, sp. nov.**

FIGS 12–15

FUNGAL NAME FN 570075




Differs from *Septobasidium pittospori* by its transverse layer at the pillar bases.

TYPE: China, Guangxi, Longzhou, Nonggang Natural Reserve, alt. 400 m, on unidentified plant, associated with tribe *Lepidosaphidini* (*Diaspididae*), 21.VII.2012, S.H. He GX02 (HMAS 244429, holotype).

ETYMOLOGY: The epithet refers to the transverse layer running below the pillars in section.

Basidiomata on branches, resupinate, 14–28 cm long, 4–6 cm wide, pale cinnamon-brown, cinnamon-brown or brown; surface smooth, peeling off at

FIGS 9–11. *Septobasidium guangxiense* (HMAS 244542, holotype). 9. Basidiomata on branches; 10. Section of basidioma; 11. Basidium (arrow). FIGS 12–14. *Septobasidium transversum* (HMAS 244429, holotype). 12. Basidiomata on branches; 13. Section of basidioma; 14. Basidia.






FIG 15. *Septobasidium transversum* (HMAS 244429, holotype). Basidia.

maturity, emerging pillars; margin determinate or indeterminate. In section 800–1200 μm thick. Subiculum brown, 30–100 μm thick. Pillars brown, 300–730 μm high, 50–190 μm wide, forming a transverse layer 20–60 μm thick at the pillar bases. Hyphal layer brown, 150–250 μm thick. Hymenium hyaline or brown, 70–120 μm thick. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or curved, 4-celled, 42–60 $\times$ 9–12 μm , hyaline or brown. Sterigmata conical, hyaline or brown. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

COMMENTS: *Septobasidium transversum* is similar to *S. pittospori* C.X. Lu & L. Guo, which differs by having shorter pillars (190–290 μm), thicker hyphal layer (340–580 μm) in the section, and without a transverse layer at the base of pillars (Lu & Guo 2011).

Including the three new species reported in this paper, 57 *Septobasidium* species have now been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a,b,c, 2010a,b,c, 2011, Lu et al. 2010, Chen & Guo 2011a,b,c, 2012a,b,c, Li & Guo 2013a,b, Li et al. 2013).

Acknowledgements

The authors would like to express their deep thanks to Drs. Eric H.C. McKenzie (Auckland, New Zealand) and Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewers, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, to Dr. Shuanghui He and Mr. Fan He for collecting specimens, to Mr. Ziyu Cao (Institute of Botany, Chinese Academy of Sciences) for identifying the

host plant, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insects, and to Mrs. Xiangfei Zhu for inking in line drawings. This study was supported by the National Natural Science Foundation of China (No. 30670005).

Literature cited

- Chen SZ, Guo L. 2011a. *Septobasidium sichuanense* sp. nov. (*Septobasidiaceae*) from China. *Mycotaxon* 115: 481–484. <http://dx.doi.org/10.5248/115.481>
- Chen SZ, Guo L. 2011b. *Septobasidium atalantiae* sp. nov. (*Septobasidiaceae*) and *S. henningsii* new to China. *Mycotaxon* 117: 291–296. <http://dx.doi.org/10.5248/117.291>
- Chen SZ, Guo L. 2011c. *Septobasidium glycosmidis* and *S. albiziae* spp. nov. (*Septobasidiaceae*) from Hainan Province. *Mycosystema* 30: 861–864.
- Chen SZ, Guo L. 2012a. Three new species and three new Chinese records of *Septobasidium* (*Septobasidiaceae*). *Mycosystema* 31: 651–655.
- Chen SZ, Guo L. 2012b. Three new species of *Septobasidium* (*Septobasidiaceae*) from Hainan Province in China. *Mycotaxon* 120: 269–276. <http://dx.doi.org/10.5248/120.269>
- Chen SZ, Guo L. 2012c. *Septobasidium saurauiae* sp. nov. (*Septobasidiaceae*) and *S. pseudopedicellatum* new to China. *Mycotaxon* 118: 283–288. <http://dx.doi.org/10.5248/118.283>
- Chen SZ, Guo L. 2013. Three new species of *Septobasidium* (*Septobasidiaceae*) from southern and southwestern China. *Mycotaxon* 121: 375–383. <http://dx.doi.org/10.5248/121.375>
- Couch JN. 1938. The genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.
- Kirschner R, Chen CJ. 2007. New reports of two hypophyllous *Septobasidium* species from Taiwan. *Fung. Sci.* 22(1,2): 39–46.
- Li W, Guo L. 2013a. Two new species of *Septobasidium* (*Septobasidiaceae*) from Yunnan Province in China. *Mycotaxon* 125: 91–96. <http://dx.doi.org/10.5248/125.91>
- Li W, Guo L. 2013b. *Septobasidium diaspidioti* sp. nov. from Anhui Province. *J. Fung. Res.* 11(4): 239–241.
- Li W, Chen SZ, Guo L, Ye YQ. 2013. *Septobasidium hoveniae* sp. nov. and *S. rhabarbarinum* new to China. *Mycotaxon* 125: 97–101. <http://dx.doi.org/10.5248/125.97>
- Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. *Mycotaxon* 109: 103–106. <http://dx.doi.org/10.5248/109.103>
- Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. *Mycotaxon* 109: 477–482. <http://dx.doi.org/10.5248/109.477>
- Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *Septobasidium kameii* new to China. *Mycotaxon* 110: 239–245. <http://dx.doi.org/10.5248/110.239>
- Lu CX, Guo L. 2010a. Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. *Mycotaxon* 112: 143–151. <http://dx.doi.org/10.5248/112.143>
- Lu CX, Guo L. 2010b. Two new species of *Septobasidium* (*Septobasidiaceae*) and *S. pallidum* new to China. *Mycotaxon* 113: 87–93. <http://dx.doi.org/10.5248/113.87>
- Lu CX, Guo L. 2010c. Two new species of *Septobasidium* (*Septobasidiaceae*) from Hainan province in China. *Mycotaxon* 114: 217–223. <http://dx.doi.org/10.5248/114.217>
- Lu CX, Guo L. 2011. Two new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. *Mycotaxon* 116: 395–400. <http://dx.doi.org/10.5248/116.395>
- Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. *Mycotaxon* 111: 269–274. <http://dx.doi.org/10.5248/111.269>
- Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.
- Tai FL. 1979. *Sylloge Fungorum Sinicorum*. Science Press, Beijing. 1527 p.
- Teng SC. 1963. *Fungi of China*. Science Press, Beijing. 808 p.